

University of Missouri, St. Louis

IRL @ UMSL

Chemistry & Biochemistry Faculty Works

Chemistry and Biochemistry

April 2011

Dicarbonyldichloridobis(Trimethylphosphane)Iron(Ii)–Carbonyldichloridotris(1/1/2)

Nigam Rath

University of Missouri–St. Louis

Meghan Stouffer

University of Washington

Matthew Janssen

University of Washington

John Bleeke

University of Washington

Follow this and additional works at: <https://irl.umsl.edu/chemistry-faculty>



Part of the [Biochemistry Commons](#), [Chemistry Commons](#), and the [Computer Sciences Commons](#)

Recommended Citation

Rath, Nigam; Stouffer, Meghan; Janssen, Matthew; and Bleeke, John,

"Dicarbonyldichloridobis(Trimethylphosphane)Iron(Ii)–Carbonyldichloridotris(Trimethylphosphane)Iron(Ii)–Tetrahydrofuran (1/1/2)" (2011). *Chemistry & Biochemistry Faculty Works*. 21.

DOI: <https://doi.org/10.1107/S1600536811009305>

Available at: <https://irl.umsl.edu/chemistry-faculty/21>

This Article is brought to you for free and open access by the Chemistry and Biochemistry at IRL @ UMSL. It has been accepted for inclusion in Chemistry & Biochemistry Faculty Works by an authorized administrator of IRL @ UMSL. For more information, please contact marvinh@umsl.edu.

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Dicarbonyldichloridobis(trimethylphosphane)iron(II)–carbonyldichlorido-tris(trimethylphosphane)iron(II)–tetrahydrofuran (1/1/2)

Nigam P. Rath,^{a*} Meghan Stouffer,^b Matthew K. Janssen^b and John R. Bleeker^b

^aDepartment of Chemistry and Biochemistry and Center for Nanoscience, University of Missouri–St Louis, 1 University Boulevard, St Louis, MO 63121-4400, USA, and

^bDepartment of Chemistry, Washington University, One Brookings Drive, St Louis, MO 63130-4899, USA

Correspondence e-mail: rathn@umsl.edu


Received 25 February 2011; accepted 10 March 2011

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}–\text{C}) = 0.004$ Å; disorder in solvent or counterion; R factor = 0.034; wR factor = 0.081; data-to-parameter ratio = 22.5.

The asymmetric unit of the title crystal, $[\text{FeCl}_2(\text{C}_3\text{H}_9\text{P})_3(\text{CO})] \cdot [\text{FeCl}_2(\text{C}_3\text{H}_9\text{P})_2(\text{CO})_2] \cdot 2\text{C}_4\text{H}_8\text{O}$, contains half molecules of the two closely related Fe^{II} complexes lying on mirror planes and a tetrahydrofuran solvent molecule, one C atom of which is disordered over two sets of sites with site occupancy factors 0.633 (9) and 0.367 (9). In both Fe^{II} complex molecules, a distorted octahedral coordination geometry has been observed around the Fe atoms. Weak intermolecular $\text{C}–\text{H} \cdots \text{O}$ interactions are observed in the crystal structure.

Related literature

For the synthetic background, see: Harris *et al.* (1978). For the crystal structure of a related complex, see: Venturi *et al.* (2004).



Experimental

Crystal data

$[\text{FeCl}_2(\text{C}_3\text{H}_9\text{P})_3(\text{CO})] \cdot [\text{FeCl}_2(\text{C}_3\text{H}_9\text{P})_2(\text{CO})_2] \cdot 2\text{C}_4\text{H}_8\text{O}$

$M_r = 862.10$
Orthorhombic, $Pnma$

$a = 10.8391$ (9) Å
 $b = 16.9670$ (12) Å
 $c = 22.2871$ (18) Å
 $V = 4098.8$ (6) Å³
 $Z = 4$

Mo $K\alpha$ radiation
 $\mu = 1.20$ mm⁻¹
 $T = 100$ K
 $0.20 \times 0.12 \times 0.10$ mm

Data collection

Bruker APEXII CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2008)
 $T_{\text{min}} = 0.800$, $T_{\text{max}} = 0.895$

144514 measured reflections
4904 independent reflections
3939 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.090$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$
 $wR(F^2) = 0.081$
 $S = 1.06$
4904 reflections
218 parameters

1 restraint
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.68$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.70$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D–H \cdots A$	$D–H$	$H \cdots A$	$D \cdots A$	$D–H \cdots A$
$\text{C42}–\text{H42B} \cdots \text{O15}^{\text{i}}$	0.98	2.53	3.422 (3)	151
$\text{C43}–\text{H43C} \cdots \text{O1}^{\text{i}}$	0.98	2.58	3.510 (3)	158
$\text{C43}–\text{H43A} \cdots \text{O1}^{\text{ii}}$	0.98	2.43	3.392 (3)	167

Symmetry codes: (i) $x - \frac{1}{2}, y, -z + \frac{3}{2}$; (ii) $-x + \frac{3}{2}, -y + 1, z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2010); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008) and *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2009).

The authors acknowledge financial support from the National Science Foundation (grant No. CHE 0420497) for the purchase of the X-ray diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2394).

References

- Bruker (2008). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
Bruker (2009). *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Bruker (2010). *APEX2*. Bruker AXS Inc., Madison, Wisconsin, USA.
Harris, T. V., Rathke, J. W. & Muettterties, E. L. (1978). *J. Am. Chem. Soc.* **100**, 6966–6977.
Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
Venturi, C., Bellachioma, G., Cardaci, G. & Macchioni, A. (2004). *Inorg. Chim. Acta*, **357**, 3712–3720.

supporting information

Acta Cryst. (2011). E67, m462 [doi:10.1107/S1600536811009305]

Dicarbonyldichloridobis(trimethylphosphane)iron(II)–carbonyldichloridotris(trimethylphosphane)iron(II)–tetrahydrofuran (1/1/2)

Nigam P. Rath, Meghan Stouffer, Matthew K. Janssen and John R. Bleeke

S1. Comment

An interesting cocrystallization has occurred from a reaction of CO with $\text{Cl}_2\text{Fe}(\text{PMe}_3)_2$, resulting in compound (I), $\text{C}_8\text{H}_{18}\text{Cl}_2\text{FeO}_2\text{P}_2$, from the addition of two equivalents of CO to $\text{Cl}_2\text{Fe}(\text{PMe}_3)_2$ and compound (II), $\text{C}_{10}\text{H}_{27}\text{Cl}_2\text{FeOP}_3$, probably from the addition of one equivalent of CO, followed by the rapid addition of one equivalent of free PMe_3 , which is present in the reaction solution. In this paper, we report the crystal structure of the two compounds, (I) and (II) which have been cocrystallized along with a molecule of tetrahydrofuran solvate per molecule of complex (Fig. 1).

The asymmetric unit of the title crystal contains half molecules of the two compounds, (I) and (II), lying on mirror planes and a molecule of tetrahydrofuran solvate, $\text{C}_4\text{H}_8\text{O}$; a carbon atom of the solvent molecule is disordered over two sites C4S and C4S' with site occupancy factors 0.633 (9) and 0.367 (9). In compound (I), the PMe_3 ligands occupying axial positions, are *trans* with respect to each other with an angle of $175.20(4)^\circ$ and the CO and Cl are *trans* with respect to each other at equatorial positions. In compound (II), the *trans* PMe_3 ligands are located at $166.41(4)^\circ$ to each other; the 3rd PMe_3 is *trans* to a Cl. The octahedral coordination is completed with the 2nd Cl being *trans* to a CO ligand. In both compounds, the ligands around Fe lie in slightly distorted octahedral coordination geometry. An overlay plot of the two molecules drawn by *Mercury* (Macrae *et al.*, 2008) shows the close similarity of the two molecules (Fig. 2).


There are weak intermolecular interactions of the type $\text{C}-\text{H}\cdots\text{O}$ which are observed between both the carbonyl O atoms of (I) and a methyl hydrogen atom of (II). The O of the solvent THF also has weak interactions with a methyl hydrogen atoms of (II) (Table 1).

S2. Experimental

FeCl_2 (0.21 g, 1.62×10^{-3} mol) and PMe_3 (0.40 ml, 3.86×10^{-3}) were stirred in 20 ml of THF for 10 min, producing a clear gray solution of $\text{Cl}_2\text{Fe}(\text{PMe}_3)_2$ (Harris *et al.*, 1978) in the presence of excess PMe_3 . Carbon monoxide was then bubbled through the solution until the color changed to an intense orange. The THF solvent was removed under vacuum and the resulting powder was extracted with pentane. After filtration through Celite, the pentane was removed under vacuum. The product was dissolved in a 1:2 mixture of THF and pentane and cooled to 243 K, causing orange crystals to form overnight.

S3. Refinement

H atoms bonded to the C atoms located on the mirror planes were located in a difference map and refined using a riding model. Other H atoms were calculated with idealized geometries with $\text{C}-\text{H} = 0.98$ and 0.99 \AA for methyl and methylene type H-atoms, respectively, and refined using a riding model with $U_{\text{iso}}(\text{H}) = 1.2$ (1.5 for methyl groups) times $U_{\text{eq}}(\text{C})$. A molecule of THF was located in the asymmetric unit wherein C4 was disordered with partial occupancy factors 0.633 (9) and 0.367 (9).

**Figure 1**

The molecular structure of (I) and (II) with atom labels and 50% probability displacement ellipsoids for non-H atoms.

Disordered atoms in the solvent are omitted for clarity. Symmetry codes represented by A in atomic labels: for (I) = $x, 0.5 - y, z$ and for (II) = $x, 1.5 - y, z$.




Figure 2

Overlay plot of the two molecules.




Figure 3

A unit cell packing plot of the title crystal; H atoms have been omitted for clarity.

Dicarbonyldichloridobis(trimethylphosphane)iron(II)– carbonyldichloridotris(trimethylphosphane)iron(II)– tetrahydrofuran (1/1/2)

Crystal data



$M_r = 862.10$

Orthorhombic, *Pnma*

$a = 10.8391(9) \text{ \AA}$

$b = 16.9670(12) \text{ \AA}$

$c = 22.2871(18) \text{ \AA}$

$V = 4098.8(6) \text{ \AA}^3$

$Z = 4$

$F(000) = 1808$

$D_x = 1.397 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 9975 reflections

$\theta = 2.6\text{--}27.4^\circ$

$\mu = 1.20 \text{ mm}^{-1}$

$T = 100 \text{ K}$

Plate, light yellow

$0.20 \times 0.12 \times 0.10 \text{ mm}$

Data collection

Bruker APEXII CCD
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: $8.3333 \text{ pixels mm}^{-1}$

φ and ω scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2008)

$T_{\min} = 0.800$, $T_{\max} = 0.895$

144514 measured reflections

4904 independent reflections

3939 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.090$

$\theta_{\max} = 27.6^\circ$, $\theta_{\min} = 1.5^\circ$

$h = -14 \rightarrow 14$

$k = -21 \rightarrow 22$

$l = -29 \rightarrow 29$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.034$
 $wR(F^2) = 0.081$
 $S = 1.06$
 4904 reflections
 218 parameters
 1 restraint
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0323P)^2 + 5.017P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.002$
 $\Delta\rho_{\max} = 0.68 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.70 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger. *SHELX* restraints used:
 delu o2 c2

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Fe1	0.74441 (4)	0.2500	0.609986 (19)	0.01347 (10)	
Cl1	0.64447 (5)	0.14941 (3)	0.66198 (2)	0.02064 (12)	
P1	0.89145 (8)	0.2500	0.68214 (4)	0.01903 (18)	
P2	0.58528 (8)	0.2500	0.54398 (4)	0.01575 (17)	
O1	0.86416 (17)	0.37737 (10)	0.54685 (8)	0.0268 (4)	
C1	0.8188 (2)	0.32598 (13)	0.57122 (10)	0.0180 (5)	
C11	0.9949 (3)	0.33405 (17)	0.68026 (13)	0.0353 (7)	
H11A	0.9465	0.3828	0.6812	0.053*	
H11B	1.0441	0.3324	0.6434	0.053*	
H11C	1.0498	0.3323	0.7152	0.053*	
C12	0.8343 (4)	0.2500	0.75831 (15)	0.0277 (8)	
H12A	0.7861	0.2036	0.7650	0.042*	
H12B	0.9084	0.2500	0.7829	0.042*	
C21	0.5823 (2)	0.33398 (15)	0.49372 (12)	0.0258 (5)	
H21A	0.5116	0.3294	0.4665	0.039*	
H21B	0.6587	0.3354	0.4703	0.039*	
H21C	0.5749	0.3826	0.5172	0.039*	
C22	0.4336 (3)	0.2500	0.57645 (16)	0.0263 (8)	
H22A	0.3697	0.2500	0.5460	0.039*	
H22B	0.4224	0.2044	0.6013	0.039*	
Fe2	0.39182 (4)	0.7500	0.91114 (2)	0.01534 (11)	
Cl2	0.17671 (7)	0.7500	0.90064 (4)	0.02288 (17)	
Cl3	0.38648 (9)	0.7500	1.01832 (4)	0.0306 (2)	

P3	0.43591 (8)	0.7500	0.81372 (4)	0.01842 (18)	
P4	0.36847 (6)	0.61635 (3)	0.91539 (3)	0.01807 (13)	
O2	0.6473 (3)	0.7500	0.92770 (12)	0.0318 (6)	
C2	0.5567 (5)	0.7500	0.92171 (15)	0.0287 (9)	
C31	0.3099 (3)	0.7500	0.76165 (16)	0.0280 (8)	
H31A	0.3440	0.7500	0.7207	0.042*	
H31B	0.2601	0.7974	0.7654	0.042*	
C32	0.5302 (2)	0.66852 (14)	0.78642 (11)	0.0229 (5)	
H32A	0.4848	0.6189	0.7912	0.034*	
H32B	0.6071	0.6661	0.8095	0.034*	
H32C	0.5493	0.6767	0.7439	0.034*	
C41	0.2881 (3)	0.58171 (15)	0.98174 (11)	0.0291 (6)	
H41A	0.2086	0.6088	0.9851	0.044*	
H41B	0.3381	0.5929	1.0174	0.044*	
H41C	0.2743	0.5248	0.9786	0.044*	
C42	0.2761 (2)	0.56933 (14)	0.85754 (11)	0.0247 (5)	
H42A	0.2698	0.5128	0.8659	0.037*	
H42B	0.3152	0.5772	0.8184	0.037*	
H42C	0.1933	0.5926	0.8572	0.037*	
C43	0.5079 (2)	0.55672 (14)	0.91735 (11)	0.0240 (5)	
H43A	0.5573	0.5712	0.9525	0.036*	
H43B	0.5560	0.5659	0.8808	0.036*	
H43C	0.4855	0.5009	0.9198	0.036*	
C1S	0.7599 (3)	0.48598 (17)	0.79495 (13)	0.0420 (7)	
H1S1	0.7804	0.4575	0.7575	0.050*	
H1S2	0.6691	0.4908	0.7978	0.050*	
C2S	0.8096 (3)	0.44219 (17)	0.84835 (14)	0.0413 (7)	
H2S1	0.7417	0.4181	0.8717	0.050*	
H2S2	0.8672	0.4001	0.8355	0.050*	
C3S	0.8759 (3)	0.50294 (18)	0.88488 (13)	0.0400 (7)	
H3S1	0.8205	0.5280	0.9146	0.048*	0.633 (9)
H3S2	0.9481	0.4801	0.9059	0.048*	0.633 (9)
C4S	0.9139 (6)	0.5590 (4)	0.8379 (3)	0.0371 (13)	0.633 (9)
H4S1	0.9280	0.6118	0.8555	0.044*	0.633 (9)
H4S2	0.9912	0.5411	0.8186	0.044*	0.633 (9)
O1S	0.8150 (2)	0.56183 (12)	0.79471 (9)	0.0454 (6)	
H3S3	0.9638	0.4884	0.8897	0.048*	0.367 (9)
H3S4	0.8382	0.5075	0.9252	0.048*	0.367 (9)
C4S'	0.8641 (11)	0.5827 (7)	0.8499 (5)	0.0371 (13)	0.367 (9)
H4S3	0.8085	0.6195	0.8713	0.044*	0.367 (9)
H4S4	0.9457	0.6080	0.8450	0.044*	0.367 (9)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Fe1	0.0166 (2)	0.0124 (2)	0.0114 (2)	0.000	-0.00049 (17)	0.000
Cl1	0.0240 (3)	0.0195 (3)	0.0184 (3)	-0.0034 (2)	0.0011 (2)	0.0051 (2)
P1	0.0192 (4)	0.0221 (4)	0.0157 (4)	0.000	-0.0032 (3)	0.000

P2	0.0188 (4)	0.0154 (4)	0.0130 (4)	0.000	-0.0015 (3)	0.000
O1	0.0345 (10)	0.0197 (9)	0.0262 (9)	-0.0062 (7)	0.0047 (8)	0.0025 (7)
C1	0.0199 (11)	0.0182 (11)	0.0159 (11)	0.0019 (9)	-0.0024 (9)	-0.0033 (9)
C11	0.0331 (15)	0.0431 (16)	0.0295 (14)	-0.0166 (12)	-0.0103 (12)	0.0045 (12)
C12	0.032 (2)	0.036 (2)	0.0151 (17)	0.000	-0.0034 (15)	0.000
C21	0.0278 (13)	0.0243 (13)	0.0253 (13)	0.0000 (10)	-0.0057 (10)	0.0076 (10)
C22	0.0202 (17)	0.035 (2)	0.0233 (18)	0.000	-0.0009 (14)	0.000
Fe2	0.0170 (2)	0.0139 (2)	0.0151 (2)	0.000	0.00190 (18)	0.000
Cl2	0.0173 (4)	0.0231 (4)	0.0282 (4)	0.000	0.0019 (3)	0.000
Cl3	0.0299 (5)	0.0303 (5)	0.0314 (5)	0.000	0.0029 (4)	0.000
P3	0.0199 (4)	0.0185 (4)	0.0168 (4)	0.000	-0.0012 (3)	0.000
P4	0.0214 (3)	0.0147 (3)	0.0181 (3)	-0.0008 (2)	0.0026 (2)	0.0005 (2)
O2	0.0430 (17)	0.0244 (14)	0.0281 (15)	0.000	0.0089 (13)	0.000
C2	0.066 (3)	0.0088 (15)	0.0116 (16)	0.000	0.0014 (18)	0.000
C31	0.0295 (19)	0.0314 (19)	0.0231 (18)	0.000	-0.0049 (15)	0.000
C32	0.0262 (12)	0.0209 (12)	0.0216 (12)	0.0000 (9)	0.0044 (10)	-0.0014 (9)
C41	0.0387 (15)	0.0201 (12)	0.0284 (13)	-0.0003 (11)	0.0112 (12)	0.0053 (10)
C42	0.0257 (13)	0.0197 (12)	0.0287 (13)	-0.0063 (10)	-0.0027 (10)	-0.0017 (10)
C43	0.0267 (13)	0.0182 (12)	0.0272 (13)	0.0027 (10)	-0.0013 (10)	-0.0005 (10)
C1S	0.066 (2)	0.0282 (15)	0.0321 (15)	-0.0039 (14)	-0.0086 (15)	-0.0017 (12)
C2S	0.055 (2)	0.0303 (15)	0.0383 (17)	-0.0008 (14)	-0.0094 (15)	0.0058 (13)
C3S	0.0549 (19)	0.0383 (17)	0.0269 (14)	-0.0017 (14)	-0.0064 (14)	0.0052 (12)
C4S	0.040 (4)	0.042 (3)	0.029 (3)	-0.013 (2)	-0.008 (2)	0.005 (2)
O1S	0.0659 (15)	0.0324 (11)	0.0378 (12)	-0.0139 (10)	-0.0191 (11)	0.0110 (9)
C3S'	0.0549 (19)	0.0383 (17)	0.0269 (14)	-0.0017 (14)	-0.0064 (14)	0.0052 (12)
C4S'	0.040 (4)	0.042 (3)	0.029 (3)	-0.013 (2)	-0.008 (2)	0.005 (2)
O1S'	0.0659 (15)	0.0324 (11)	0.0378 (12)	-0.0139 (10)	-0.0191 (11)	0.0110 (9)

Geometric parameters (Å, °)

Fe1—C1 ⁱ	1.749 (2)	P4—C42	1.817 (2)
Fe1—C1	1.749 (2)	P4—C43	1.819 (2)
Fe1—P1	2.2641 (10)	O2—C2	0.991 (5)
Fe1—P2	2.2670 (9)	C31—H31A	0.9837
Fe1—C11 ⁱ	2.3300 (6)	C31—H31B	0.9732
Fe1—C11	2.3300 (6)	C32—H32A	0.9800
P1—C12	1.807 (4)	C32—H32B	0.9800
P1—C11	1.814 (3)	C32—H32C	0.9800
P1—C11 ⁱ	1.814 (3)	C41—H41A	0.9800
P2—C22	1.796 (4)	C41—H41B	0.9800
P2—C21	1.813 (2)	C41—H41C	0.9800
P2—C21 ⁱ	1.813 (2)	C42—H42A	0.9800
O1—C1	1.139 (3)	C42—H42B	0.9800
C11—H11A	0.9800	C42—H42C	0.9800
C11—H11B	0.9800	C43—H43A	0.9800
C11—H11C	0.9800	C43—H43B	0.9800
C12—H12A	0.9565	C43—H43C	0.9800
C12—H12B	0.9731	C1S—O1S	1.419 (3)

C21—H21A	0.9800	C1S—C2S	1.503 (4)
C21—H21B	0.9800	C1S—H1S1	0.9899
C21—H21C	0.9800	C1S—H1S2	0.9896
C22—H22A	0.9693	C2S—C3S	1.497 (4)
C22—H22B	0.9597	C2S—H2S1	0.9895
Fe2—C2	1.803 (5)	C2S—H2S2	0.9908
Fe2—P3	2.2232 (10)	C3S—C4S	1.472 (6)
Fe2—P4 ⁱⁱ	2.2837 (6)	C3S—H3S1	0.9900
Fe2—P4	2.2838 (6)	C3S—H3S2	0.9900
Fe2—Cl2	2.3433 (9)	C4S—O1S	1.442 (6)
Fe2—Cl3	2.3895 (11)	C4S—H4S1	0.9900
P3—C31	1.792 (4)	C4S—H4S2	0.9900
P3—C32	1.823 (2)	C4S'—H4S3	0.9900
P3—C32 ⁱⁱ	1.823 (2)	C4S'—H4S4	0.9900
P4—C41	1.814 (2)		
C1 ⁱ —Fe1—C1	94.97 (15)	C31—P3—Fe2	117.95 (13)
C1 ⁱ —Fe1—P1	91.50 (8)	C32—P3—Fe2	116.50 (8)
C1—Fe1—P1	91.50 (8)	C32 ⁱⁱ —P3—Fe2	116.50 (8)
C1 ⁱ —Fe1—P2	91.74 (8)	C41—P4—C42	99.88 (13)
C1—Fe1—P2	91.74 (8)	C41—P4—C43	101.48 (12)
P1—Fe1—P2	175.20 (4)	C42—P4—C43	103.34 (12)
C1 ⁱ —Fe1—Cl1 ⁱ	179.61 (8)	C41—P4—Fe2	114.12 (9)
C1—Fe1—Cl1 ⁱ	85.42 (7)	C42—P4—Fe2	117.88 (8)
P1—Fe1—Cl1 ⁱ	88.51 (2)	C43—P4—Fe2	117.46 (8)
P2—Fe1—Cl1 ⁱ	88.22 (2)	O2—C2—Fe2	179.8 (4)
C1 ⁱ —Fe1—Cl1	85.42 (7)	P3—C31—H31A	108.3
C1—Fe1—Cl1	179.61 (8)	P3—C31—H31B	111.6
P1—Fe1—Cl1	88.51 (2)	H31A—C31—H31B	106.7
P2—Fe1—Cl1	88.22 (2)	P3—C32—H32A	109.5
Cl1 ⁱ —Fe1—Cl1	94.19 (3)	P3—C32—H32B	109.5
C12—P1—C11	103.51 (12)	H32A—C32—H32B	109.5
C12—P1—C11 ⁱ	103.51 (12)	P3—C32—H32C	109.5
C11—P1—C11 ⁱ	103.6 (2)	H32A—C32—H32C	109.5
C12—P1—Fe1	115.20 (13)	H32B—C32—H32C	109.5
C11—P1—Fe1	114.74 (9)	P4—C41—H41A	109.5
C11 ⁱ —P1—Fe1	114.74 (9)	P4—C41—H41B	109.5
C22—P2—C21	103.45 (11)	H41A—C41—H41B	109.5
C22—P2—C21 ⁱ	103.45 (11)	P4—C41—H41C	109.5
C21—P2—C21 ⁱ	103.62 (18)	H41A—C41—H41C	109.5
C22—P2—Fe1	115.78 (12)	H41B—C41—H41C	109.5
C21—P2—Fe1	114.50 (9)	P4—C42—H42A	109.5
C21 ⁱ —P2—Fe1	114.50 (9)	P4—C42—H42B	109.5
O1—C1—Fe1	177.4 (2)	H42A—C42—H42B	109.5
P1—C11—H11A	109.5	P4—C42—H42C	109.5
P1—C11—H11B	109.5	H42A—C42—H42C	109.5
H11A—C11—H11B	109.5	H42B—C42—H42C	109.5
P1—C11—H11C	109.5	P4—C43—H43A	109.5

H11A—C11—H11C	109.5	P4—C43—H43B	109.5
H11B—C11—H11C	109.5	H43A—C43—H43B	109.5
P1—C12—H12A	109.5	P4—C43—H43C	109.5
P1—C12—H12B	104.3	H43A—C43—H43C	109.5
H12A—C12—H12B	111.3	H43B—C43—H43C	109.5
P2—C21—H21A	109.5	O1S—C1S—C2S	107.5 (2)
P2—C21—H21B	109.5	O1S—C1S—H1S1	110.2
H21A—C21—H21B	109.5	C2S—C1S—H1S1	110.2
P2—C21—H21C	109.5	O1S—C1S—H1S2	110.1
H21A—C21—H21C	109.5	C2S—C1S—H1S2	110.3
H21B—C21—H21C	109.5	H1S1—C1S—H1S2	108.5
P2—C22—H22A	111.9	C3S—C2S—C1S	105.2 (2)
P2—C22—H22B	110.4	C3S—C2S—H2S1	110.8
H22A—C22—H22B	108.2	C1S—C2S—H2S1	110.7
C2—Fe2—P3	85.09 (11)	C3S—C2S—H2S2	110.5
C2—Fe2—P4 ⁱⁱ	96.00 (2)	C1S—C2S—H2S2	110.7
P3—Fe2—P4 ⁱⁱ	93.689 (19)	H2S1—C2S—H2S2	108.9
C2—Fe2—P4	96.00 (2)	C4S—C3S—C2S	101.1 (3)
P3—Fe2—P4	93.687 (19)	C4S—C3S—H3S1	111.6
P4 ⁱⁱ —Fe2—P4	166.40 (4)	C2S—C3S—H3S1	111.6
C2—Fe2—Cl2	178.23 (11)	C4S—C3S—H3S2	111.6
P3—Fe2—Cl2	96.68 (4)	C2S—C3S—H3S2	111.6
P4 ⁱⁱ —Fe2—Cl2	83.905 (19)	H3S1—C3S—H3S2	109.4
P4—Fe2—Cl2	83.906 (19)	O1S—C4S—C3S	106.8 (4)
C2—Fe2—Cl3	83.88 (11)	O1S—C4S—H4S1	110.4
P3—Fe2—Cl3	168.98 (4)	C3S—C4S—H4S1	110.4
P4 ⁱⁱ —Fe2—Cl3	87.469 (19)	O1S—C4S—H4S2	110.4
P4—Fe2—Cl3	87.471 (19)	C3S—C4S—H4S2	110.4
Cl2—Fe2—Cl3	94.34 (3)	H4S1—C4S—H4S2	108.6
C31—P3—C32	102.17 (11)	C1S—O1S—C4S	106.3 (3)
C31—P3—C32 ⁱⁱ	102.17 (11)	H4S3—C4S'—H4S4	108.9
C32—P3—C32 ⁱⁱ	98.60 (16)		
C1 ⁱ —Fe1—P1—C12	-132.49 (7)	P4 ⁱⁱ —Fe2—P3—C32	153.62 (10)
C1—Fe1—P1—C12	132.49 (7)	P4—Fe2—P3—C32	-37.81 (10)
Cl1 ⁱ —Fe1—P1—C12	47.118 (16)	Cl2—Fe2—P3—C32	-122.09 (9)
Cl1—Fe1—P1—C12	-47.117 (16)	Cl3—Fe2—P3—C32	57.91 (9)
C1 ⁱ —Fe1—P1—C11	107.43 (14)	C2—Fe2—P3—C32 ⁱⁱ	-57.91 (9)
C1—Fe1—P1—C11	12.42 (14)	P4 ⁱⁱ —Fe2—P3—C32 ⁱⁱ	37.81 (10)
Cl1 ⁱ —Fe1—P1—C11	-72.95 (12)	P4—Fe2—P3—C32 ⁱⁱ	-153.62 (9)
Cl1—Fe1—P1—C11	-167.19 (12)	Cl2—Fe2—P3—C32 ⁱⁱ	122.09 (9)
C1 ⁱ —Fe1—P1—C11 ⁱ	-12.42 (14)	Cl3—Fe2—P3—C32 ⁱⁱ	-57.91 (9)
C1—Fe1—P1—C11 ⁱ	-107.43 (14)	C2—Fe2—P4—C41	111.10 (15)
Cl1 ⁱ —Fe1—P1—C11 ⁱ	167.19 (12)	P3—Fe2—P4—C41	-163.46 (11)
Cl1—Fe1—P1—C11 ⁱ	72.96 (12)	P4 ⁱⁱ —Fe2—P4—C41	-40.7 (2)
C1 ⁱ —Fe1—P2—C22	132.49 (7)	Cl2—Fe2—P4—C41	-67.12 (11)
C1—Fe1—P2—C22	-132.49 (7)	Cl3—Fe2—P4—C41	27.52 (11)
Cl1 ⁱ —Fe1—P2—C22	-47.128 (16)	C2—Fe2—P4—C42	-132.18 (15)

C11—Fe1—P2—C22	47.126 (16)	P3—Fe2—P4—C42	-46.74 (10)
C1 ⁱ —Fe1—P2—C21	-107.25 (12)	P4 ⁱⁱ —Fe2—P4—C42	76.00 (19)
C1—Fe1—P2—C21	-12.23 (12)	C12—Fe2—P4—C42	49.60 (10)
C11 ⁱ —Fe1—P2—C21	73.13 (10)	C13—Fe2—P4—C42	144.24 (10)
C11—Fe1—P2—C21	167.38 (10)	C2—Fe2—P4—C43	-7.49 (15)
C1 ⁱ —Fe1—P2—C21 ⁱ	12.23 (12)	P3—Fe2—P4—C43	77.96 (10)
C1—Fe1—P2—C21 ⁱ	107.25 (12)	P4 ⁱⁱ —Fe2—P4—C43	-159.30 (17)
C11 ⁱ —Fe1—P2—C21 ⁱ	-167.39 (10)	C12—Fe2—P4—C43	174.30 (10)
C11—Fe1—P2—C21 ⁱ	-73.14 (10)	C13—Fe2—P4—C43	-91.06 (10)
C2—Fe2—P3—C31	180.0	O1S—C1S—C2S—C3S	11.0 (4)
P4 ⁱⁱ —Fe2—P3—C31	-84.29 (2)	C1S—C2S—C3S—C4S	-28.5 (4)
P4—Fe2—P3—C31	84.29 (2)	C2S—C3S—C4S—O1S	36.8 (5)
C12—Fe2—P3—C31	0.0	C2S—C1S—O1S—C4S	11.8 (4)
C13—Fe2—P3—C31	180.000 (1)	C3S—C4S—O1S—C1S	-31.1 (6)
C2—Fe2—P3—C32	57.91 (9)		

Symmetry codes: (i) $x, -y+1/2, z$; (ii) $x, -y+3/2, z$.

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C42—H42B \cdots O1S ⁱⁱⁱ	0.98	2.53	3.422 (3)	151
C43—H43C \cdots O1 ⁱⁱⁱ	0.98	2.58	3.510 (3)	158
C43—H43A \cdots O1 ^{iv}	0.98	2.43	3.392 (3)	167

Symmetry codes: (iii) $x-1/2, y, -z+3/2$; (iv) $-x+3/2, -y+1, z+1/2$.