

7-11-2018

# Alternatives in Magnesium Sulfate Treatment Courses: An Investigation of Physician Prescribing Behaviors

Erin Williams  
elwpf4@mail.umsl.edu

Follow this and additional works at: <https://irl.umsl.edu/dissertation>



Part of the [Maternal, Child Health and Neonatal Nursing Commons](#)

---

## Recommended Citation

Williams, Erin, "Alternatives in Magnesium Sulfate Treatment Courses: An Investigation of Physician Prescribing Behaviors" (2018). *Dissertations*. 782.  
<https://irl.umsl.edu/dissertation/782>

This Dissertation is brought to you for free and open access by the UMSL Graduate Works at IRL @ UMSL. It has been accepted for inclusion in Dissertations by an authorized administrator of IRL @ UMSL. For more information, please contact [marvinh@umsl.edu](mailto:marvinh@umsl.edu).

Alternatives in Magnesium Sulfate Treatment Courses: An Investigation of Physician  
Prescribing Behaviors

---

Doctor of Nursing Practice Project Presented to the  
Faculty of Graduate Studies  
University of Missouri – St. Louis

---

In Partial Fulfillment of the Requirements  
For the Degree of Doctor of Nursing Practice

by

Erin L. Williams, BSN, RNC-MNN

---

DNP Committee Chair: Nancy Magnuson, DNS, CS, FNP-BC, RN

DNP Committee Member: Roberta Lavin, Ph.D., FNP-BC, FAAN

DNP Committee Member: Roxanne Vandermause, PhD, RN

DNP Committee Member: Heidi Tymkew, PT, DPT, MHS, CSS

Graduation August 2018

### Abstract

**Problem:** The gold standard for managing preeclampsia in the postpartum patient is 24-hours of magnesium sulfate as this is the time the patient is most susceptible to experiencing a seizure (American College of Obstetricians & Gynecologists, 2013). However, these patients are at an increased risk for experiencing adverse health outcomes, as well as interruptions in maternal-infant bonding due to the mother requiring a higher level of care.

**Methods:** A one group pretest-posttest design was conducted to gain a better understanding of the current practice while also assessing the willingness of physicians to change prescribing behavior after receiving the research-based education.

**Results:** During the study period, 49 obstetric providers agreed to participate in the study. There was no statistically significant difference between survey results regarding likelihood to prescribe a reduced course of magnesium sulfate ( $p=.821$ ). However, there was a statistically significant difference seen between the perception that 24-hours of magnesium sulfate is necessary to manage preeclampsia ( $p=.020$ ) and the theory that patients with preeclampsia without severe features could benefit from an abbreviated treatment course ( $p=.012$ ).

**Implications for Practice:** While data results do not indicate an immediate change in the prescribing of magnesium sulfate, physicians expressed interest in continued efforts to study this topic. There is an opportunity to improve patient safety by standardizing patient education. Additionally, a discrepancy was noted between the physicians' perceptions of the postpartum acuity of the preeclamptic patient. Bridging the gap between nursing and

providers ensures that providers have a reasonable expectation of the postpartum experience.

### **Introduction**

Preeclampsia is a leading cause of maternal morbidity and preterm labor contributing to 9%-26% of life-threatening maternal outcomes globally (Townsend, O'Brien, Khalil, 2016, p. 79). The American College of Obstetricians and Gynecologists (2013) defines preeclampsia as a blood pressure higher than 140/90 on two separate accounts at least four hours apart in a previously normotensive woman after 20 weeks' gestation and proteinuria (p. 4). Preeclampsia can also be diagnosed with hypertension after 20 weeks' gestation in the absence of proteinuria, but with signs of multiple organ involvement resulting in thrombocytopenia, renal insufficiency, and impaired liver function (ACOG, 2013, p. 4). The American College of Obstetricians and Gynecologists (2013) has identified magnesium sulfate infusions to be the gold standard in preventing seizures in patients with preeclampsia (2013, p. 6). However, prolonged durations of magnesium sulfate is shown to increase the risk of blood clots due to increased time to ambulation, increased risk of urinary tract infections related to the duration of indwelling catheter, and interrupting of maternal-infant bonding due to the mother needing higher-level care resulting in the delay in breastfeeding (Maia, Katz, Neto, Caiado, Azevedo & Amorim, 2014, p. 260). Traditionally, preeclampsia is treated with a magnesium sulfate infusion for the first 24-hours after delivery since that is the period the patient is most susceptible to experiencing seizures related to high blood pressures (Maia et al., 2014, p. 260). However, current literature suggests that a 12-hour treatment course of magnesium sulfate can be as practical as a 24-hour treatment course in preventing seizures in low-risk preeclamptic patients (Anjum, Goel, Sharma, Mohsin, & Garg, 2016, p. 68; Kashanian,

Koohpayehzadeh, Sheikhsari, Barapour, Sahraian, & Asadolla, 2016, p. 2282; Maia et al., 2014, p. 260).

This project took place within the obstetric (OB) department on a sizeable academic campus located in the city. The majority of patients cared for on these units come from low-income, underserved areas of St. Louis (Barnes-Jewish Hospital, 2017).

Demographically, this community is primarily black accounting for 96.8% of its residents (Statistical Atlas, 2015). There is data to support that when compared to Caucasians, African Americans are at higher risk of experiencing preeclampsia (Breathett, Muhlestein, & Gulati, 2014, p. 886). In addition to race other risk factors include “women who are ...primigravida, have multiparous pregnancy, prior history of preeclampsia, aged 35 years or older, and those who have existing chronic hypertension, diabetes, chronic kidney disease, obesity, or connective tissue disorder” (Breathett, Muhlestein, & Gulati, 2014, p. 887).

Anjum et al. (2016) acknowledge the financial impact this treatment has on not only the medical facility but also the patient (p. 71). Knowing the financial strains faced by this patient population a change in the protocol can result in benefits for the patient financially, as well as physiologically. Prescribing an abbreviated treatment course of magnesium sulfate could result in effectively preventing seizures in low-risk postpartum patients with preeclampsia while also improving clinical outcomes, increasing patient satisfaction, and reducing treatment costs. Recognizing that changing magnesium treatment time from 24-hours to 12-hours would be a significant change in practice, assessing the physicians' willingness to change prescriptive behavior will be a critical first step to what could develop into a larger research project for the future. The purpose

of this project is to determine whether implementation of a research-based educational session for OB physicians, to discuss new knowledge related to abbreviated magnesium sulfate treatment of preeclampsia in postpartum patients, results in the participating OB physicians' willingness to change the current practice of prescribing magnesium sulfate infusion for a 24-hour period to an abbreviated 12-hour period in preeclamptic postpartum patients.

### **Review of Literature**

A thorough database search was conducted including PubMed, CINAHL, and Scopus. The search was limited to the five-year period from 2012-2017, to reflect the most current and relevant literature. Search words included magnesium sulfate, randomized controlled trials, meta-analysis, preeclampsia, postpartum, infusions, drug administration schedule, puerperal disorders, treatment outcome, postnatal care, postpartum care, bed rest, and duration. Inclusion criteria included randomized controlled trials. Additionally, articles that were not peer-reviewed, scholarly articles with nonpertinent titles and abstracts, as well as articles that were not available in full-text were excluded.

The primary concern was postpartum patients with preeclampsia treated with magnesium sulfate remaining on strict bed rest. On the postpartum unit at Barnes-Jewish Hospital, it was once expected that these patients remain on strict bed rest for multiple reasons. Having preeclampsia predisposes patients to become eclamptic, or having seizures. Patients with preeclampsia are kept on seizure precautions while on a magnesium sulfate infusion requiring oxygen and suction to be readily available. Furthermore, magnesium sulfate acts as a smooth muscle relaxer placing these patients at

risk for falls (Murray & McKinney, 2010, pp. 637-638). As of recently, physicians were beginning to allow patients on magnesium sulfate to leave the patient care unit without nurse supervision, and without oxygen and suction readily available. Unfortunately, there was no literature available to support the need for strict bed rest for patients treated with magnesium sulfate which resulted in a change in search strategies to address this problem from a different perspective. Further concerns for patients treated with magnesium sulfate is the length of time with a Foley catheter, which increases the risk for urinary tract infections, the period to ambulation, and length of time to maternal-infant bonding (Maia et al., 2014, p. 260). Utilizing these critical concerns as search strategies introduced the topic of abbreviated treatment courses of magnesium sulfate in preventing seizures in postpartum women with preeclampsia.

There is currently no standard recommendation for the length of time a patient with preeclampsia should be treated with magnesium sulfate to prevent seizures (Anjum et al., 2016a, p. 68). However, preeclampsia is traditionally treated for the first 24-hours after delivery because that is the period the patient is most at risk for experiencing (Anjum et al., 2016a, p. 68; Darngawn, Jose, Regi, Bansal, & Jeyaseelan, 2012, p. 237; Maia et al., 2014, p. 260). Throughout the literature, several approaches were taken regarding reducing treatment time. Chama, Geidam, Bako, Mairiga & Atterwahmie (2013) was unique in that the preferred method of administration was by intramuscular injection (IM) and utilized the Pritchard method (p. 131). The Pritchard method involves the patient receiving an initial loading dose of 4 grams of magnesium sulfate by intravenous (IV) infusion along with a 5-gram magnesium sulfate IM injection into each buttock then followed by an additional 5-gram magnesium sulfate IM injection in each



buttock, alternating sides, every four hours until the 24-hour mark is reached (Chama et al., 2013, p. 132). Chama et al. (2013) argued that a shortened course of magnesium sulfate would be as efficient as the standard Pritchard method at preventing seizures (p. 131). A reduced course included the loading dose of 10 mg IV magnesium sulfate followed by only two magnesium sulfate IM injections into each buttock four hours apart (Chama et al., 2013, p. 132). It was determined that this shortened treatment course was as useful as the Pritchard method in preventing seizures in the first 24-hours after delivery ( $p=0.684$ ) while also exhibiting similar “maternal and perinatal outcomes” and reducing the total dose of magnesium sulfate by more than 40%” (Chama et al., 2013, p. 133).

Previous research has shown that the loading dose of magnesium sulfate alone could be useful in preventing seizures in the first 24-hours after delivery (Anjum et al., 2016a, p. 68; Anjum, Rajaram, & Bano, 2016, p. 983). However, a theme of reducing treatment time to 6-hours was more common. Anjum et al. (2016b) compared patients receiving the traditional 24-hour treatment course to patients who received an abbreviated 6-hour treatment course of magnesium sulfate. Study groups were found to be comparable with no statistically significant differences in demographic or clinical characteristics (Anjum et al., 2016b, p. 984). Moreover, neither group exhibited seizures after completion of treatment (Anjum et al., 2016, p. 984). There was no difference between study groups concerning perinatal or neonatal outcomes (Anjum et al., 2016, p. 985). However, statistically significant differences were seen in the intervention group regarding maternal outcomes as illustrated by decreases in the dose of magnesium sulfate given Group A  $15.1 \pm 5.4$ , Group B  $42.3 \pm 7.3$ ;  $p<0.001$ ) time with a Foley catheter

(Group A  $11.3 \pm 5.1$ , Group B  $38.3 \pm 7.3$ ;  $p < 0.001$ ), and hours of monitoring (Group A  $11.1 \pm 4.7$ , Group B  $38.4 \pm 7.2$ ;  $p < 0.001$ ) (Anjum et al., 2016, p. 985). Darngawn, Jose, Bansal & Jevaseelan (2012) did similar work in reducing treatment time from 24-hours to 6-hours (p. 237). The intervention group experienced a reduction in magnesium sulfate administration, as well as reduced mean time spent receiving nursing care (23.00 versus 70.13 minutes,  $P < 0.001$ ) (Darngawn et al., 2012, p. 237-238). Vigil-De Garcia, Ramirez, Duran & Quintero (2017) also tested a 6-hour treatment course; however, their study required that patients receive less than 8-hours of magnesium sulfate treatment before delivery (p. 1). Comparable to other studies examining the 6-hour treatment time no differences were found between the study groups regarding the incidence of eclampsia (Vigil-De Garcia et al., 2017, p. 3). It was also suggested that a minimum dose of 14-grams of magnesium sulfate versus the 32-gram dose administered during the traditional 24-hour treatment is as effective in preventing seizures (Vigil-De Garcia, 2017, p. 5).

In addition to exploring the benefits of a 6-hour treatment course, preceding scholars have examined the effectiveness of a less extreme reduction in treatment time to 12-hours. Anjum et al. (2016a) compared study groups who received the traditional 24-hour treatment course of magnesium sulfate and the abbreviated 12-hour course (p. 68). Researchers found that there was no statistically significant difference between study groups regarding clinical and demographic characteristics (Anjum et al., 2016a, p. 69). Neither group exhibited seizures or toxicity after the completion of treatment; however, the control group receiving the traditional 24-hours of magnesium sulfate experienced ten seizures after two hours of therapy (Anjum et al., 2016a, p. 69). However, statistically significant differences were seen between the two groups when considering the

secondary outcomes including length of hospital stay (Study group:  $5.3 \pm 0.8$  days, Control group:  $7.5 \pm 1.5$  days;  $P < 0.001$ ) and days with a Foley catheter Study group:  $19.6 \pm 2.5$  hours, Control group:  $31.5 \pm 3.2$  hours,  $P < 0.001$ ) (Anjum et al., 2016a, p. 70). The group who received the 24-hour treatment course was shown to have received a higher dose of magnesium sulfate (Study group:  $23.2 \pm 2.8$  grams; Control group:  $34.9 \pm 3.2$  grams;  $P < 0.001$ ) when compared to the intervention group (Anjum et al., 2016a, p. 70).

Maia et al. (2014) also examined a 12-hour treatment course to the 24-hour treatment course and observed similar results. All study participants received an IV infusion of magnesium sulfate at 1-gram per hour for the first 12-hours after delivery (Maia et al., 2014, p. 261). Once the initial 12-hours had passed the infusion was discontinued for participants in the intervention group while the control group continued to receive the magnesium sulfate for the full 24-hours (Maia et al., 2014, p. 261). Similar to preceding studies, the reduced treatment time of 12-hours was as effective in preventing seizures during the first 24-hours after delivery (Maia et al., 2014, p. 261). Additionally, the magnesium sulfate did not need to be restarted for the incidence of eclampsia (Maia et al., 2014, p. 261). Along with participant treated adequately for the prevention of seizures, there was also a substantial reduction in the time participants had a Foley catheter (14.3 hours vs 25.3 hours;  $P < 0.001$ ), time to ambulation (18.8 hours vs 25.8 hours;  $P < 0.001$ ), and time to bond with baby (29.6 hours vs 35.0 hours;  $p = 0.03$ ) (Maia et al., 2014, p. 263). With the reduction in time to have contact with the baby, there could be an improvement in the “likelihood of establishing breastfeeding” (Maia et al., 2014, p. 263).

Kashanian et al. (2016) performed a similar study in investigating the effectiveness of a shortened treatment time of magnesium sulfate for 12-hours as compared to 24-hours for patients with severe preeclampsia (p. 2282). Baseline demographic and clinical characteristics were similar between both groups (Kashanian et al., 2016, p. 2283). Laboratory values including AST, ALT, alkaline phosphatase, hemoglobin, platelet count, BUN, and creatinine before and after treatment were compared between both groups and no significant difference was observed (Kashanian et al., 2016, p. 2283). In addition to comparing laboratory values, researchers also compared both groups for the incidence of side effects often associated with magnesium sulfate treatment and the transition to eclampsia including “flushing, nausea and vomiting, head lightness, headache, blurred vision, and epigastric pain” (Kashanian et al., 2016, p. 2286). There was no significant difference seen between the two groups concerning the adverse effects (Kashanian et al., 2016, p. 2286).

El-Khayat, Atef, Abdelatty, & El-senary (2016) compared all three methods, the loading dose alone, the abbreviated 12-hour course and the 24-hour course, in their capability of preventing seizures in patients with preeclampsia (p. 154). All three groups exhibited no statistically significant difference in baseline demographic and clinical characteristics (El-Khayat et al., 2016, p. 155). There was also no statistically significant difference in blood pressure, the incidence of hemorrhage, and type of delivery (El-Khayat et al., 2016, p. 155). Additionally, there were no statistically significant differences seen in neonatal outcomes between the three groups, including “prematurity, IUGR, perinatal death and NICU admission (El-Khayat et al., 2016, p. 155). In considering the primary outcome of eclampsia, there was no statistically significant

difference found between the three groups (El-Khayat et al., 2016, p. 155). There was, however, statistically significant differences noted between the three groups when considering the incidence of flushing, an adverse effect of magnesium sulfate treatment (El-Khayat et al., 2016, p. 155). Researchers concluded that following a protocol that requires a *loading dose only* would require “less maternal monitoring for urine output and blood pressure” (El-Khayat et al., 2016, p. 158).

Many of these studies shared comparable primary outcome measures that focused on the incidence of seizure activity or the recurrence of seizures. However, there was more variability seen in the secondary outcome measures which put the spotlight on clinical characteristics such as laboratory findings, participant recovery, maternal and neonatal outcomes, reports of pain, and time spent on monitoring by healthcare staff. The strength of these studies lies in the study design. Each study discussed was a randomized controlled trial. While most were single-blinded, El-Kayat et al. (2016) managed to maintain a double-blinded investigation adding to its strength (p. 154). The primary limitation of research investigating the effectiveness of an abbreviated treatment course of magnesium sulfate in preventing seizures in patients with preeclampsia is the sample size. Sample sizes of the previously mentioned studies ranged from 98 participants to 284 participants. Maia et al. (2014) recognized that larger sample sizes would be needed to observe the frequency of eclampsia in patients with a reduced treatment time of magnesium sulfate (p. 263). Specifically, the involvement of multiple facilities and upward of 18,000 participants would be necessary to have sufficient power to make a recommendation for practice change (Maia et al., 2014, p.263). Another limitation of these studies was the lack of measurement tools involved to standardize reportable data.

Darngawn et al. (2012) was the only study that made mention of an established measurement tool, the Wong-Bake pain scale, to standardize the participants' reports of pain (p. 238). Maia et al. (2014) utilized a Likert scale to measure the participant's satisfaction with the care she receiving (p. 261). All other studies relied merely on self-report from participants.

The Johns Hopkins Nursing Evidence-Based Practice Model (JHNEBP) will be utilized to implement this project. The JHNEBP Model relies on “practice, education, and research” as its pillars for professional nursing (Dearholt & Dang, 2012, p. 34). “The 18-step JHNEBP process occurs in 3 phases and can be simply described as Practice question, Evidence, and Translation (PET)” (Dearholt & Dang, 2012, p. 42). The JHNEBP Model walks the professional nurse through the process of implementing change within their organization, “provides tools for process and critique, including question development, evidence rating scale, and research and non-research evidence appraisal” (Schaffer, Sandau, & Diedrick, 2013).

## **Method**

### **Design**

The overall approach of this project will be that of change implementation. The primary goal will not be implementing the change itself but assessing the willingness of physicians to change current practice. A quality improvement design was utilized to evaluate the effectiveness of the intervention. Each group of physicians was presented the same survey, and no randomization occurred. A one group pretest-posttest design was conducted to determine if the research-based education had an impact on the physicians' willingness to change current practice.

**Setting**

This project took place on a sizable academic medical campus located in the city, as well as the offices of private physician groups. Conference rooms were utilized as arranged with administrative assistants and the physicians themselves.

**Sample**

A convenience sample of obstetric physicians that care for patients at Barnes-Jewish Hospital served as the sample population. The experience level of the physicians ranged from residents, year one through year four, fellows, attending physicians, and nurse practitioners. There are roughly 32 obstetric residents. In addition to the residents, there are two laborists, four Women's Health Nurse Practitioners (WHNP) and several groups of private attending physicians. The goal was to have at least 30 providers in the sample population.

**Approval Processes**

Since project work took place at Barnes-Jewish Hospital, a project proposal was submitted to the Department of Research's Protocol Review Committee. Because the proposed study methods do not include Barnes-Jewish Hospital team members who fall under the purview of Patient Care Services, it was not necessary to participate in the Proposal Review Committee process. Additionally, an application was submitted through the University of Missouri-St. Louis Internal Review Board and the study was deemed exempt. Risks to participants included slight discomfort if personal views were challenged in the group discussion. Ethical considerations were centered around protecting the confidentiality and anonymity of participants. While there is always a risk of breach of confidentiality, every effort to preserve privacy was maintained. No names

were written on the surveys and only identified by a prewritten number. Care was taken to ensure that responses remained confidential by having physicians place their responses in blank envelopes. Participation was voluntary and implied consents acknowledged with the receipt of the surveys.

### **Procedures**

Providers caring for obstetric patients at Barnes-Jewish Hospital were enrolled between March 22, 2018, and May 22, 2018. A physician champion was identified to help organize appropriate meeting times, as well as ensure the educational material was suitable for presenting to the providers. Before the educational sessions, packets were made for each participant that included the implied consent form, a brief synopsis of the evidence as well as the surveys taken before and after the presentation. An unspecified number of meetings were planned with each session not lasting longer than 30 minutes. Providers were contacted to schedule time for the primary investigator to present the research-based educational sessions where evidence that supports the possible change in practice from treating preeclampsia with 24-hours of magnesium sulfate to reduced time of 12-hours was presented. Each provider received an information sheet, and their survey completion implied their understanding of the research and consent to participate (see Appendix C). Physicians were asked to complete a questionnaire before and after the presentation. Physicians were not required to include their name on the surveys, and they were left in an unmarked envelope to preserve anonymity and left in a specified location.

### **Data Collection/Analysis**

Self-reported questionnaires were administered to providers before the session and then immediately after to determine if the research-based educational session had an



impact on the providers' willingness to change the current prescribing of magnesium sulfate treatment courses. Both the pre-test survey and the post-test survey had nine questions which were identical. Both pre-test and post-test survey's asked the physicians to identify what level provider they are with choices being resident, fellow, attending, or nurse practitioner. The questionnaires were brief and focused on the providers prior knowledge of magnesium sulfate treatment courses, if the session introduced new information, beliefs on various concepts of care of the preeclamptic patient, and if the evidence presented encouraged them to change prescribing behaviors (see Appendices A and B for examples of surveys). A paired t-test using SPSS version 24 (IBM, Armonk, NY, USA) was anticipated to be utilized to determine if there is a statistically significant difference between the willingness to change practice before and after the intervention. Furthermore, a Chi-square test was utilized to determine if any association existed between provider level and change in variables.

A manual chart review of patients receiving magnesium sulfate between December 15, 2016, through December 21, 2018 was conducted with the assistance of the clinical pharmacist. Extracted data from a hospital computer included the date of the prescribed medication to note the frequency of prescription, the level of the provider prescribing the medication to gain an understanding of who is most frequently ordering the medication and the duration of the treatment. Because special privileges were necessary to access Centricity GE, additional data to determine the duration of treatment was unable to be obtained. Identifiable data was not utilized or recorded. Data was entered and analyzed on a spreadsheet. That list was used to determine the dates of treatment to identify the frequency of magnesium sulfate ordering for preeclampsia as well as the provider level.

### **Results**

A manual chart review of physician orders of magnesium sulfate for postpartum preeclampsia resulted in a total of 811 orders, which averages 16 orders per week. Of those 811 orders, 93% of them were entered by OB residents with the majority of the ordering being completed by second-year and fourth-year residents. The remaining 7% of orders represented fellows, nurse practitioners, laborists, service attending physicians, and private service attending physicians. The OB residents are primarily responsible for entering orders for patients' which explains the high percentage of magnesium sulfate orders completed by residents. The majority of preeclampsia diagnoses are made on the antepartum unit where second-year residents reside, while the fourth-year residents are chief residents who primarily manage the treatment plans on all obstetric patients. Understanding of the residents' rotation helps to explain the distribution of medication orders.

In the present quality improvement project, a convenience sample of 49 providers was surveyed. Among those 49 providers, 26 were residents, 5 were fellows, 15 were attendings, and 3 were nurse practitioners (see appendix D). A paired-samples t-test was used to determine whether there was a statistically significant mean difference between survey answers after participants listened to an educational presentation (see Appendix E). Statistically significant differences were observed regarding the disruptive role a 24-hour course of magnesium sulfate plays in breastfeeding and maternal-infant bonding ( $M=.878$ ,  $SD=1.130$ ,  $t=5.437$ ,  $p=.000$ ) and the necessity of the patient to have a Foley catheter allowing nursing to adequately monitor for magnesium sulfate toxicity ( $M=.571$ ,

SD=1.258,  $t=3.179$ ,  $p=.003$ ). While there was a change in the perception that patients should be kept on strict bedrest while receiving this high-risk medication, it was not statistically significant ( $M=-.102$ ,  $SD=.368$ ,  $t=-1.943$ ,  $p=.058$ ). A Fisher's Exact test was conducted between provider level and the belief that a patient should remain on bedrest for the duration of treatment. There was no statistically significant association between provider level and idea that a patient should stay on bedrest for the duration of treatment, ( $p=.066$ ) (see Appendix F, Figure 1).

There was a significant difference in the scores for the necessity for patients to receive a full 24-hours of magnesium sulfate ( $M=-.347$ ,  $SD=1.011$ ,  $t=-2.401$ ,  $p=.020$ ) and if patients with stable preeclampsia could benefit from an abbreviated magnesium sulfate course ( $M=.449$ ,  $SD=1.209$ ,  $t=2.600$ ,  $p=.012$ ). While providers appeared to be more accepting of an abbreviated treatment course of magnesium sulfate in managing patients with preeclampsia, there was not a statistically significant difference seen in the providers' willingness to prescribe a 12-hour course of magnesium sulfate ( $M=0.41$ ,  $SD=1.258$ ,  $t=.227$ ,  $p=.821$ ). Because the mean difference was not statistically significant, we can reject the alternative hypothesis and accept the null hypothesis.

A chi-square test for association was conducted to determine if there were any differences in willingness to change prescribing behavior between residents and attendings. The data was analyzed for both pretest and posttest surveys. A response of "unsure" was considered to be a "no" for this analysis. All expected cell frequencies were greater than five for pretest results. There was no statistically significant association between provider level and willingness to prescribe a reduced treatment course on the pretest,  $\chi^2(1) = .360$ ,  $p = .548$  (see Appendix F, figure 2). Posttest results exhibited an

expected cell frequency of four in which case a Fisher's Exact test was conducted. Again, there was not a statistically significant association noted between provider level and willingness to prescribe a reduced treatment course,  $p = .064$  (see Appendix F, figure 3). While attending physician responses remained consistent pretest and posttest, an unexpected increase in the number of resident physicians unwilling to change their prescriptive behavior was observed posttest. These results suggest that while the education presented had an impact on the providers ability to understand various implications experienced by patients receiving 24-hours of magnesium sulfate, ultimately providers are not yet committed to prescribing reduced treatment courses of magnesium sulfate (see Appendix G).

### **Discussion**

This study revealed that there was no statistically significant difference in the willingness of the providers to prescribe a reduced treatment course of magnesium sulfate for postpartum patients with preeclampsia ( $p=.821$ ). However, there was a statistically significant difference noted between the providers concerning the belief that postpartum magnesium sulfate could be reduced from the traditional 24-hour treatment course ( $p=.020$ ). Providers also responded well to the notion that patients that present with preeclampsia without severe features could, in fact, benefit from an abbreviated treatment course ( $p=.012$ ). This lack of correlation highlights that while providers are open to the concept of a reduced treatment course, they are not yet committed to a change in treatment plan.

This study began with the primary investigator's concern for patient safety while they were being allowed out of bed and off the hospital unit while receiving this high-risk

medication infusion. Additionally, these patients were lacking a Foley catheter, which decreases the ability of the nurse to monitor for magnesium sulfate adequately. It is imperative to strictly monitor the patient's urine output to ensure that they have adequate diuresis (30 mL/hr) as oliguria (>30 mL/hr) leads to the buildup of magnesium sulfate leading to toxicity (Murray & McKinney, 2010). If a patient is allowed off the hospital unit without a Foley catheter, the nurse is unable to monitor their urine output. While there was a statistically significant difference observed in the providers belief that a patient should have a Foley catheter while being treated with magnesium sulfate ( $p=.003$ ), there was no difference noted in their belief that these patients should remain on strict bedrest until treatment is complete ( $p=.058$ ).

From discussions had with providers after the presentations, the primary investigator noticed a range of opinions regarding the management of these patients. Overall, most of the attending physicians were unaware that patients with preeclampsia treated with magnesium sulfate were being allowed off the unit with the infusion running; furthermore, they were unaware that these same patients were without a Foley catheter. Resident physicians are primarily responsible for managing the care of these patients throughout their hospital stay with the guidance of the attending physicians which could explain the lack of awareness. At the time of the completion of this study, there has been recent communication with a Maternal-Fetal Medicine (MFM) physician that there are changes to come regarding managing postpartum patients with preeclampsia receiving magnesium sulfate. Additionally, the Director of the Residency Program communicated the same sentiment to the resident physicians. Further discussion with the Washington University Obstetric Physician group highlighted that these patients requiring a higher

level of care are included in the nurses assignment of six to eight patients, including mothers and newborns. It became clear that the obstetric providers were unaware of the acuity level of the postpartum nursing assignments as they were only aware that these patients received one-to-one care on the labor and delivery unit.

An opportunity exists to standardize patient education through their journey from labor and delivery to the postpartum unit. As stated above, since these patients receive one-to-one care from the labor and delivery nurse there is more opportunity for the patient to be taken to see their baby in the neonatal intensive care unit (NICU) by their nurse, as well as receive closer monitoring. The labor and delivery staff, nursing and providers, lead these patients to expect the same level of care when they transfer to the postpartum unit without consideration to the staffing differences as well as increased acuity. As this project comes to an end work is in progress to update current nursing policies that address the management of care of patients receiving magnesium sulfate. Additionally, patient education is being standardized to ensure these patients expectations are managed and safety ensured throughout their stay. It is the plan of the primary investigator to collect additional data to determine if patients receiving magnesium sulfate present to the postpartum unit with Foley catheters and remain on strict bedrest after presenting to the obstetric providers.

### **Conclusion**

While obstetric providers seem to be open to the idea of an abbreviated treatment course of magnesium sulfate, they are not yet comfortable with changing their prescriptive behavior. Given this fact, further research is necessary to show the clinical benefits of an abbreviated treatment course of magnesium sulfate. This study highlights

that the management of care for postpartum patients with preeclampsia receiving magnesium sulfate can be improved. While not necessarily indicated by survey results, providers are now aware of the risk these patients are at for experiencing adverse health outcomes due to being allowed off the unit and not having a Foley catheter throughout treatment time.

Work to standardize patient education is being done so that staff on labor and delivery and postpartum are providing the same information to the patients to manage their expectations throughout their stay. Dissemination of staff education will occur at the annual Women & Infant's Skills Day in conjunction with a presentation by the clinical pharmacist on medication information for magnesium sulfate. Creating a partnership with the Washington University Obstetric Physicians, as well as the Maternal-Fetal Medicine Physicians, could help fill the gaps in the perception of the nursing care provided on the postpartum unit for the patients on magnesium sulfate. Follow-up is expected to occur with Maternal-Fetal Medicine as a physician with the Washington University Obstetric group found that this study would make an exceptional Fellow project. Continued efforts with Maternal-Fetal Medicine could help provide the additional research to support further a reduced treatment course of magnesium sulfate for patients receiving magnesium sulfate.

### References

- American College of Obstetricians and Gynecologists. (2013). Hypertension in Pregnancy. Retrieved from <https://www.acog.org/~media/Task%20Force%20and%20Work%20Group%20Reports/public/HypertensioninPregnancy.pdf>.
- Anjum, S., Goel, N., Sharma, R., Mohsin, Z., & Garg, N. (2016). Maternal outcomes after 12 hours and 24 hours of magnesium sulfate therapy for eclampsia. *International Journal of Gynecology and Obstetrics*, 132(1), 68-71.
- Anjum, S., Rajaram, G.P., & Bano, I. (2016). Short-course postpartum (6-h) magnesium sulfate therapy in severe preeclampsia. *Archives of Gynecology and Obstetrics*, 293(5), 983-986. Doi: 10.1007/s00404-015-3903-y.
- Barnes-Jewish Hospital. (2017). Disparities Dashboard. Retrieved from <https://www.barnesjewish.org/About-Us/Community-Benefit/Community-Health-Needs-Assessment>.
- Breathett, K., Muhlestein, D., Foraker, R., & Gulati, M. (2014). Differences in preeclampsia rates between African American and Caucasian women: Trends from the National Hospital Discharge Survey. *Journal of Women's Health*, 23(11), 886-893. Doi: 10.1089/jwm.2014.4749.
- Chama, C.M., Geidam, A.D., Bako, B., Mairiga, A.G., & Atterwahmie, A. (2013). A shortened versus standard matched postpartum magnesium sulphate regimen in the treatment of eclampsia: A randomised controlled trial. *African Journal of Reproductive Health*, 17(3), 131-136.
- City-data. (2017). Jeff-Vander-Lou neighborhood. Retrieved from <http://www.city-data.com/neighborhood/JeffVanderLou-Saint-Louis-MO.html>
- Darngawn, L., Jose, R., Regi, A., Bansal, R., & Jeyaseelan, L. (2012). A shortened postpartum magnesium sulfate prophylaxis regime in pre-eclamptic women at low risk of eclampsia. *International Journal of Gynecology & Obstetrics*, 116(3): 237-239. Doi: 10.1016/j.ijgo.2011.09.028.
- Dearholt, S.L. & Dang, D. (2012). *Johns Hopkins Nursing Evidence-Based Practice: Model and Guidelines*. (2<sup>nd</sup> ed.). Indianapolis, IN: Sigma Theta Tau International.
- El-Khayat, W., Atef, A., Abdelatty, S., & El-semari, A. (2016). A novel protocol for postpartum magnesium sulphate in severe pre-eclampsia: A randomized controlled pilot trial. *Journal of Maternal-Fetal and Newborn Medicine*, 29(1), 154-158. Doi: 10.3109/14767058.2014.991915.
- Kashanian, M., Koochpayehzadeh, J., Sheikhsari, N., Bararpour, F., Sahraian, G., & Asadolla, S. (2016). A comparison between the two methods of magnesium sulfate administration for duration of 12 versus 24 h after delivery in patients with severe preeclampsia. *The Journal of Maternal-Fetal & Neonatal Medicine*, 29(14): 2282-2287. Doi: 10.3109/14767058.2015.1083547
- Maia, S.B., Katz, L., Neto, C.N., Caiado, B.V.R., Azevedo, A.P.R.L., & Amorim, M.M.R. (2014). Abbreviated (12-hour) versus traditional (24-hour) postpartum magnesium sulfate therapy in severe preeclampsia. *International Journal of Gynecology and Obstetrics*, 126(3), 260-264.
- Murray, S.S. & McKinney, E.S. (2010). *Foundations of Maternal-Newborn and Women's Health Nursing* (5<sup>th</sup> Ed.). Maryland Heights, MO: Saunders Elsevier.



- Schaffer, M.A., Sandau, K.E., & Diedrick, L. (2013). Evidence-based practice models for organizational change: Overview and practical applications. *Journal of Advanced Nursing*, 69(5), 1197-1209. Doi: 10.1111/j.1365-2648.2012.06122.x.
- Statistical Atlas. (2015). Jeffvanderlou, St. Louis, Missouri. Retrieved from <https://statisticalatlas.com/neighborhood/Missouri/St-Louis/Jeffvanderlou>.
- Townsend, R., O'Brien, P., & Khalil, A. (2016). Current best practice in the management of hypertensive disorders in pregnancy. *Integrated Blood Pressure Control*, 9, 79-94. Doi: 10.2147.IBPC.S77344.
- Vigil-De Garcia, P., Ramirez, R., Duran, Y. & Quintero, A. (2017). Magnesium sulfate for 6 vs 24 hours post-delivery in patients who received magnesium sulfate for less than 8 hours before birth: a randomized clinical trial. *BMC Pregnancy and Birth*, 17(1): 241-246. doi: 10.1186/s12884-017-1424-3.

**Appendix A**

**Postpartum Magnesium Sulfate Prescribing Pre-Survey**

Physician Level (please circle)				
Resident	Fellow	Attending	Nurse Practitioner	
<b>1. Are you aware of the current evidence that suggests that for some patients a reduced treatment time could be as effective as the traditional 24-hour course?</b>				
Yes		No		
<b>2. Should patients get out of bed while receiving magnesium sulfate infusions?</b>				
Yes		No		
<b>3. Patients receiving magnesium sulfate are at risk for experiencing a hemorrhage.</b>				
1 <small>Strongly unlikely</small>	2	3	4	5 <small>Strongly likely</small>
<b>4. Treatment with magnesium sulfate should follow a 24-hour treatment course</b>				
1 <small>Strongly agree</small>	2	3	4	5 <small>Strongly disagree</small>
<b>5. Patients presenting with stable preeclampsia could benefit from an abbreviated magnesium sulfate treatment course</b>				
1 <small>Strongly agree</small>	2	3	4	5 <small>Strongly disagree</small>
<b>6. Patients on magnesium sulfate must have an indwelling catheter in place to ensure accurate urine output</b>				
1 <small>Strongly agree</small>	2	3	4	5 <small>Strongly disagree</small>
<b>7. Breastfeeding and maternal-infant bonding are interrupted for patients being treated with magnesium sulfate</b>				
1 <small>Strongly agree</small>	2	3	4	5 <small>Strongly disagree</small>
<b>8. Patients receiving magnesium sulfate are at risk for falls.</b>				
1 <small>Strongly agree</small>	2	3	4	5 <small>Strongly disagree</small>
<b>9. How likely are you to prescribe a reduced treatment time of magnesium sulfate (12-hours)</b>				
1 <small>Strongly unlikely</small>	2	3	4	5 <small>Strongly likely</small>

**Appendix B**

**Postpartum Magnesium Sulfate Prescribing Post-Survey**

Physician Level (please circle)				
Resident	Fellow	Attending	Nurse Practitioner	
1. Are you aware of the current evidence that suggests that for some patients a reduced treatment time could be as effective as the traditional 24-hour course?				
Yes		No		
2. Should patients get out of bed while receiving magnesium sulfate infusions?				
Yes		No		
3. Patients receiving magnesium sulfate are at risk for experiencing a hemorrhage.				
1 Strongly unlikely	2	3	4	5 Strongly likely
4. Treatment with magnesium sulfate should follow a 24-hour treatment course.				
1 Strongly agree	2	3	4	5 Strongly disagree
5. Patients presenting with stable preeclampsia could benefit from an abbreviated magnesium sulfate treatment course.				
1 Strongly agree	2	3	4	5 Strongly disagree
6. Patients on magnesium sulfate must have an indwelling catheter in place to ensure accurate urine output.				
1 Strongly agree	2	3	4	5 Strongly disagree
7. Breastfeeding and maternal-infant bonding are interrupted for patients being treated with magnesium sulfate.				
1 Strongly agree	2	3	4	5 Strongly disagree
8. Patients receiving magnesium sulfate are at risk for falls.				
1 Strongly agree	2	3	4	5 Strongly disagree
9. How likely are you to prescribe a reduced treatment time of magnesium sulfate (12-hours)				
1 Strongly unlikely	2	3	4	5 Strongly likely

Appendix C



**College of Nursing**

8001 Natural Bridge Road  
 St. Louis, Missouri 63121-4499  
 E-mail: elwpf4@umsl.edu

**Informed Consent for Participation in Research Activities**

Alternatives in Magnesium Sulfate Treatment Courses: An Investigation of Physician Prescribing Behaviors

HSC Approval Number \_\_\_\_\_

Principal Investigator \_\_\_\_\_ Erin Williams, BSN, RNC-MNN \_\_\_\_\_

PI's Phone Number 314-223-7929

**Why am I being asked to participate?**

You are invited to participate in a research study about exploring the benefits of alternative treatment courses for magnesium sulfate and physician willingness to change prescribing behavior conducted by Erin Williams, graduate student at the University of Missouri-St. Louis and Barnes-Jewish Hospital. You have been asked to participate in this research because you are a practicing obstetric physician and may be eligible to participate. We ask that you read this form and ask any questions you may have before agreeing to be in the research. Your participation in this research is voluntary. Your decision whether to participate will not affect your current or future relations with the University or Barnes-Jewish Hospital. If you decide to participate, you are free to withdraw at any time without affecting that relationship.

**What is the purpose of this research?**

The purpose of this research is to examine alternative treatment courses of magnesium sulfate infusions for low-risk patients with preeclampsia and determine the willingness of physicians to change prescribing behavior.

**What procedures are involved?**

If you agree to participate in this research, you can expect:

- You will be asked to answer questions on a survey, listen to an educational presentation, and then answer questions on another survey.
- It will take about 35 minutes to complete the survey, listen to the presentation, and answer the second survey.

Approximately 50 participants may be involved in this research at Barnes-Jewish Hospital.

**What are the potential risks and discomforts?**

There may be a slight risk of discomfort if personal views are challenged in the group discussion. There is also risk of breach of confidentiality regarding study participation, however, every effort to preserve confidentiality will be maintained.

**Are there benefits to taking part in the research?**

There are no direct benefits for you participating in this study. However, this research will offer feedback to the researcher on how physicians perceive different treatment courses of magnesium sulfate in treating preeclampsia. The results may help the researcher understand prescriptive behavior.

**Appendix C (continued)****What about privacy and confidentiality?**

Surveys will not contain names and will be identified only by a letter at the upper left corner to maintain anonymity. Surveys will be given as a packet to each participant at the time of the educational session. The pre- and post-survey will be differentiated by color and participants will be instructed which survey to complete at the appropriate time. The surveys will be turned in as one packet to a predetermined location away from the principal investigator. By agreeing to participate, you understand and agree that your data may be shared with other researchers and educators in the form of presentations and/or publications. In all cases, your identity will not be revealed. In rare instances, a researcher's study must undergo an audit or program evaluation by an oversight agency (such as the Office for Human Research Protection). That agency would be required to maintain confidentiality of your data. In addition, all data will be stored on a password-protected computer and/or in a locked office.

**What are the costs for participating in this research?**

There is no cost for participating in this research.

**Will I be paid for my participation in this research?**

There is no form of payment for your participation in this study.

**Can I withdraw or be removed from the study?**

Your participation is voluntary and you may choose not to participate in this research study or to withdraw your consent at any time. You may choose not to answer any questions you do not want to answer. You will NOT be penalized in any way should you choose not to participate or withdrawn.

**Who should I contact if I have questions?**


The researcher conducting this study is Erin Williams. You may ask any questions you have now. If you have questions later, you may contact the researcher at 314-223-7929.

**What are my rights as a research subject?**

If you have any questions about your rights as a research subject, you may call the Chairperson of the Institutional Review Board at (314) 516-5897.

**I have read this consent form and my completion of this survey implies my understanding and consent to my participation in the research described above.**

**Appendix D**



---

**Figure 1: Sample population by provider level**

**Appendix E**

**Paired Samples Test**

		Mean	Std. Deviation	Std. Error Mean	Paired Differences		t	df	Sig. (2-tailed)
					95% Confidence Interval of the Difference				
					Lower	Upper			
Pair 1	Lit_pre - Literature_post	.224	.422	.060	.103	.346	3.728	48	.001
Pair 2	Bedrest_pre - Bedrest_post	-.102	.368	.053	-.208	.004	-1.943	48	.058
Pair 3	Hemorrhage_pre - Hemorrhage_post	-.082	.534	.076	-.235	.072	-1.071	48	.290
Pair 4	Treatment_pre - Treatment_post	-.347	1.011	.144	-.637	-.056	-2.401	48	.020
Pair 5	Stable_pre - Stable_post	.449	1.209	.173	.102	.796	2.600	48	.012
Pair 6	Catheter_pre - Catheter_post	.571	1.258	.180	.210	.933	3.179	48	.003
Pair 7	Breastfeeding_pre - Breastfeeding_post	.878	1.130	.161	.553	1.202	5.437	48	.000
Pair 9	Falls_pre - Falls_post	.245	.855	.122	-.001	.490	2.006	48	.051
Pair 10	Reduced_pre - Reduced_post	.041	1.258	.180	-.320	.402	.227	48	.821

**Table 1: Paired t-test results for survey responses. *P* values less than .05 were considered significant.**

**Appendix F**

**Chi-Square Tests**

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	3.776 <sup>a</sup>	1	.052		
Continuity Correction <sup>b</sup>	2.422	1	.120		
Likelihood Ratio	4.331	1	.037		
Fisher's Exact Test				.066	.055
Linear-by-Linear Association	3.684	1	.055		
N of Valid Cases	41				

a. 1 cells (25.0%) have expected count less than 5. The minimum expected count is 3.51.

b. Computed only for a 2x2 table

**Table 1: Association between provider level and belief that patients' should remain on bedrest**

**Chi-Square Tests**

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.360 <sup>a</sup>	1	.548		
Continuity Correction <sup>b</sup>	.067	1	.796		
Likelihood Ratio	.357	1	.550		
Fisher's Exact Test				.734	.395
Linear-by-Linear Association	.352	1	.553		
N of Valid Cases	41				

a. 0 cells (.0%) have expected count less than 5. The minimum expected count is 5.12.

b. Computed only for a 2x2 table

**Table 2: Pretest association between provider level and willingness to change prescribing behavior**

**Chi-Square Tests**

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	4.742 <sup>a</sup>	1	.029		
Continuity Correction <sup>b</sup>	3.282	1	.070		
Likelihood Ratio	4.635	1	.031		
Fisher's Exact Test				.064	.036
Linear-by-Linear Association	4.626	1	.031		
N of Valid Cases	41				


a. 1 cells (25.0%) have expected count less than 5. The minimum expected count is 4.02.

b. Computed only for a 2x2 table


**Table 3: Posttest association between provider level and willingness to change prescribing behavior**



**Appendix G**



**Graph 1: Provider level and willingness to change prescriptive behavior pretest**



**Graph 2: Provider level and willingness to change prescriptive behavior posttest**